

Cannabinoids in pancreatic cancer: Correlation with survival and pain

Christoph W. Michalski^{1,2}*, Florian E. Oti², Mert Erkan^{1,2}, Danguole Sauliunaite^{1,2}, Frank Bergmann³, Pal Pacher⁴, Sandor Batkai⁴, Michael W. Müller^{1,2}, Nathalia A. Giese², Helmut Friess^{1,2} and Jörg Kleeff^{1,2}

¹Department of General Surgery, Technische Universität Munich, Munich, Germany

²Department of General Surgery, University of Heidelberg, Heidelberg, Germany

³Institute of Pathology, University of Heidelberg, Heidelberg, Germany

⁴Section of Oxidative Stress Tissue Injury, Laboratory of Physiologic Studies, National Institutes of Health, NIAAA, Bethesda, MD

Cannabinoids exert antiproliferative properties in a variety of malignant tumors, including pancreatic ductal adenocarcinoma (PDAC). In our study, we quantitatively evaluated the immunoreactivity for cannabinoid-1 (CB1) and cannabinoid-2 (CB2) receptors as well as for the endocannabinoid metabolizing enzymes fatty acid amide hydrolase (FAAH) and monoacyl glycerol lipase (MGLL). Furthermore, quantitative real-time RT-PCR for CB1, CB2, FAAH and MGLL in normal pancreas and pancreatic cancer tissues was performed. Levels of endocannabinoids were determined by liquid chromatography/mass spectrometry. Immunoreactivity scores and QRT-PCR expression levels were correlated with the clinico-pathological (TNM, survival, pain) status of the patients. Evaluation of endocannabinoid levels revealed that these remained unchanged in PDAC compared to the normal pancreas. Patients with high CB1 receptor levels in enlarged nerves in PDAC had a lower combined pain score (intensity, frequency, duration; $p = 0.012$). There was a significant relationship between low CB1 receptor immunoreactivity or mRNA expression levels ($p = 0.0011$ and $p = 0.026$, respectively), or high FAAH and MGLL cancer cell immunoreactivity ($p = 0.036$ and $p = 0.017$, respectively) and longer survival of PDAC patients. These results are underlined by a significant correlation of high pain scores and increased survival ($p = 0.0343$). CB2 receptor immunoreactivity, CB2 receptor, FAAH and MGLL mRNA expression levels did not correlate with survival. Therefore, changes in the levels of endocannabinoid metabolizing enzymes and cannabinoid receptors on pancreatic cancer cells may affect prognosis and pain status of PDAC patients.

© 2007 Wiley-Liss, Inc.

Key words: pancreatic cancer; cannabinoids; pain; survival

Pancreatic ductal adenocarcinoma (PDAC) is a devastating disease which is scarcely affected by any therapeutic regimen due to its aggressive growth, early metastases and poor response to chemo-, radio- and targeted therapies.^{1,2} Severe pain and cachexia render the late, palliative stages of the disease a medical challenge for which new treatment strategies are needed.^{3,4}

In recent years, there has been increasing interest in cannabinoids due to their antineoplastic, anticachectic and analgesic potential. Synthetic cannabinoids mimic the function of their endogenous counterparts, such as anandamide (AEA) and arachidonoylglycerol (AG), by activating specific $G_{i/o}$ protein-coupled cannabinoid receptors.⁵ Of these, cannabinoid receptor-1 (CB1)⁶ is abundantly found in neural tissues, while cannabinoid receptor-2 (CB2)⁷ is particularly expressed on cells of the immune system. Endocannabinoids are then metabolized by enzymes such as fatty acid amide hydrolase (FAAH)⁸ and monoacyl glycerol lipase (MGLL).^{9,10} Growth inhibitory activities of cannabinoids have been demonstrated for various malignant tumors, including brain, breast, prostate, colorectal, skin and recently also pancreatic cancer.^{11–19} Regarding pancreatic cancer, these effects have not only been shown *in vivo*, but also in xenograft models.¹⁷ Cannabinoids can induce apoptosis in cancer cells, but have also been shown to inhibit tumor vascularization by altering blood vessel morphology and decreasing proangiogenic factors such as VEGF.¹⁹ Furthermore, they have been shown to decrease invasiveness of cancer cells.^{14,20–22}

Most recently, stress-regulated protein p8 has been identified as a downstream target of the proapoptotic sphingolipid ceramide,

serving as an essential mediator of growth inhibitory properties of cannabinoids by upregulating endoplasmic reticulum stress-related genes ATF-4, CHOP and TRB3.²³ Importantly, some of these effects are communicated by the endocannabinoid system *in vitro*, and inhibition of endocannabinoid inactivation has revealed tumor growth inhibition comparable to that observed with exogenously administered cannabinoids.²⁴ However, cannabinoids in some cases also exert growth-promoting activities through growth factor/receptor systems or by suppression of an antitumor immune response.^{13,25,26}

Altogether, antiproliferative effects seem to prevail, and therefore cannabinoids are considered potential candidate drugs in anti-tumor therapies, simultaneously alleviating cancer-related pain and cachexia. In line with a variety of *in vitro* studies and animal experiments, a recent study showed some response of malignant gliomas to intratumoral cannabinoid infusions in humans, underlining a potential prospect for their *in vivo* use.²⁷

Here, we concomitantly analyze the expression of cannabinoid receptors and endocannabinoid metabolizing enzymes. Our results demonstrate a correlation between longer survival in pancreatic cancer patients, and low CB1 receptor or high FAAH as well as MGLL levels in tissue specimens.

Material and methods

Patients and tissue sampling

Tissue samples were collected from patients after pancreatic resections for PDAC ($n = 40$). Normal pancreatic tissue samples were obtained through an organ donor procurement program whenever there was no suitable recipient for pancreas transplantation ($n = 10$). Pancreatic tissues were immediately snap frozen at -80°C or formalin fixed and paraffin embedded. The use of human tissue for the analysis was approved by the local ethical committee (University of Heidelberg, Germany), and written informed consent was obtained from the patients.

Measurement of endocannabinoids

Pancreas tissues for the measurement of endocannabinoids were immediately snap frozen in liquid nitrogen after resection and as an internal quality control, tissues with low numbers of mRNA expression levels of housekeeping genes were not used for analysis. Levels of AEA, 1-arachidonoylglycerol (1-AG) and 2-arachidonoylglycerol (2-AG) in frozen human pancreas samples (normal pancreas: $n = 11$; pancreatic cancer: $n = 5$) were determined by

The first three authors contributed equally to this paper.

Grant sponsors: Intramural Research program of National Institutes of Health/NIAAA, Heidelberg University.

*Correspondence to: Department of General Surgery, Technische Universität Munich, Ismaningerstrasse 22, D-81675 Munich, Germany. Fax: +49-89-4140-4870. E-mail: christoph.michalski@gmx.de

Received 24 June 2007; Accepted after revision 1 August 2007

DOI 10.1002/ijc.23114

Published online 17 October 2007 in Wiley InterScience (www.interscience.wiley.com).

liquid chromatography/mass spectrometry as described previously.²⁸

Immunohistochemical analysis of human pancreatic specimens

Immunohistochemistry was performed on paraffin-embedded sections of human pancreata (CB1/CB2: 10 normal pancreas, 40 pancreatic cancer tissues; FAAH/MGLL: 10 normal pancreas samples, 31 pancreatic cancer tissues) as described previously.^{29,30} Formalin fixed, paraffin embedded consecutive human pancreatic cancer tissue sections (3 μ m) were deparaffinized in Roticlear (Roth, Karlsruhe, Germany) and rehydrated in progressively decreasing concentrations of ethanol. Antigen retrieval was performed by boiling tissue sections in 10 mM citrate buffer (pH 6.0) in a microwave oven for 15 min. The activity of endogenous peroxidase was blocked with 3% hydrogen peroxide in methanol for 10 min at room temperature. After the peroxidase block, slides were washed in wash buffer (1 \times TBS/0.1%BSA, pH 7.4). Non-specific binding sites were blocked by incubating the slides with 1 \times TBS/3%BSA for 30 min at room temperature. Sections were subsequently incubated with rabbit polyclonal anti-CB1, anti-CB2, anti-FAAH and anti-MAGL antibodies (all from Cayman Chemical, Ann Arbor, MI) at 4°C over night at antibody dilutions (in 1 \times TBS/3%BSA) of 1:150, 1:300, 1:150 and 1:200, respectively. Specificity controls included pre-adsorption of the primary antibody with the corresponding blocking peptide (1:1) for 1 hr at 37°C. After incubation with the primary antibodies, slides were washed twice with wash buffer containing 0.05% Tween20 and once with plain wash buffer. Secondary antirabbit antibody, HRP-labeled polymer (DakoCytomation, ready to use; Hamburg, Germany) was applied for 45 min at room temperature. Slides were then washed twice using 1 \times TBS/0.1%BSA/0.05% Tween20 and once with plain wash buffer. The color reaction was performed applying DAB-chromogen/H₂O₂ substrate mixture (DakoCytomation, Hamburg, Germany) to tissue sections. Slides were counterstained in Mayer's hematoxylin, washed in tap water, dehydrated in progressively increasing concentrations of ethanol, cleared in Roticlear and mounted with permount (Biomedica, Foster City, CA). Analysis and semiquantitative evaluation were performed using an Axioplan-2 microscope (Carl Zeiss, Jena, Germany).

Semiquantitative evaluation of CB receptor, FAAH and MGLL levels in human pancreatic specimens

Immunoreactivity of CB1 and CB2 receptors as well as FAAH and MGLL on cancer cells and/or nerves was quantitatively evaluated according to intensity and area: the staining intensity of pancreatic cancer cells themselves was recorded as "no staining (0)", "weak to moderate staining (1)" or "strong staining (2)". The area of stained cancer cells was recorded as < 33% (1), 33–66% (2) or >66% (3) of all cancer cells. These numbers were then multiplied resulting in a score of 0–6. Regarding intrapancreatic nerves, only the staining intensity (0–2; as for cancer cells) was recorded due to the generally low number of nerves in pancreatic cancer tissue specimens.

Quantitative real-time RT-PCR

Quantitative real-time RT-PCR (QRT-PCR) was performed on a second, larger cohort of 53 pancreatic cancer samples as previously described.^{31,32} Primers for CB1-receptor, CB2-receptor, FAAH and MGLL were obtained from Search LC, Heidelberg, Germany. CB1 receptor primers flank the sequence NM_016083 at nucleotide 926–1164; CB2 receptor NM_001841 at nucleotide 507–738; FAAH NM_001441 at 1706–1965 and MGLL NM_007283 at nucleotide 1112–1400.

Statistical analysis

Statistical analysis was performed using the GraphPad Prism 4 Software (GraphPad, San Diego, CA). The Kaplan-Meier method and log-rank tests were used for survival analysis. Pain scores and

endocannabinoid levels were compared using the Mann-Whitney U test. Statistical significance was set at $p < 0.05$. Results are expressed as mean \pm SEM.

Results

High cancer cell levels of CB1 are associated with shorter survival

Immunohistochemical analyses of normal human pancreata revealed weak immunoreactivity for CB1 and CB2 in pancreatic acini, nerves, blood vessels and ductal cells (data not shown). Subsequently, CB receptor immunoreactivity on pancreatic cancer tissues was semiquantitatively evaluated and a combined score (staining intensity multiplied by the percentage of the stained area) was calculated. Tissue specimens were stratified accordingly into those with low CB receptor levels (represented by scores of 0–3) and those with high CB receptor levels (represented by scores of >3). Apart from the staining in the normal parts of the pancreatic cancer tissues, we observed CB1 and CB2 immunopositivity of stellate cells particularly in the vast areas of desmoplasia. Additionally, infiltrating mononuclear cells were immunopositive for CB2. In cancer cells of pancreatic cancer tissue samples, heterogeneous immunoreactivity was observed: whereas a low combined score of CB1 staining intensity was observed in 24 out of 37 samples, 13 out of 37 samples showed a high CB1 score. For CB2, there was a low staining score in 18 samples and a high immunoreactivity score in 19 tissues. Furthermore, cancer cells within single tissue samples showed various extents of CB1 and CB2 staining, ranging from no immunoreactivity to strong immunoreactivity (Figs. 1a–1c and 1g–1i). Consecutive sections which were incubated with the primary antibodies and the corresponding blocking peptide showed the specificity of the staining (Figs. 1d–1f and 1j–1l).

Semiquantitative analysis of staining intensities revealed a statistically significant negative correlation between the extent of CB1 immunoreactivity of pancreatic cancer cells and patient survival ($p = 0.0011$; Fig. 2a). The median survival in the group with low CB1 staining in cancer cells was 16 months, versus 6 months for the group with a high combined CB1 staining score. For CB2, however, no such correlation was observed ($p = 0.869$; median survival 10.75 versus 12.5 months; Fig. 2b). Furthermore, there was no correlation between tumor grade/TN(M) status and pain score of the patients (Spearman rho: pain score versus grade: $r = -0.25$, $p = 0.3$; pain score versus T status: $r = -0.21$, $p = 0.36$; pain score versus N status: $r = 0$, $p = 1.0$). Since all patients were resectable, none had distant metastases at the time of the operation. In accordance with the published literature on pancreatic cancer, the nodal status correlated significantly with overall survival (Spearman rho; $p = 0.016$).

Subsequently, tissue specimens from a second, larger cohort of patients ($n = 53$) were evaluated using quantitative RT-PCR. These analyses confirmed the results of the immunohistochemistry and revealed that higher CB1 mRNA expression levels (above median) were associated with a shorter survival (Fig. 2c; 10 months versus 16 months, $p = 0.026$) whereas CB2 mRNA expression levels were not related to patient survival (Fig. 2d; 13 months versus 15 months, $p = 0.195$).

Low cancer cell FAAH and MGLL levels are associated with shorter survival

Immunohistochemistry of normal human pancreata revealed weak immunoreactivity for FAAH and MGLL in pancreatic islets and nerves (data not shown). In pancreatic cancer tissues, mainly pancreatic cancer cells were immunopositive for FAAH and MGLL while the stroma was unstained. Enlarged intrapancreatic nerves were heterogeneously immunoreactive for FAAH and MGLL. Quantitative evaluation of pancreatic cancer cells in tissue specimens demonstrated highly variable FAAH and MGLL immunoreactivities (Fig. 3). Few samples (3 out of 29) were unstained




FIGURE 1 – Cannabinoid receptor immunostaining in human pancreatic cancer tissues. Low (*a, g*), moderate (*b, h*) and strong (*c, i*) immunostaining for CB1 or CB2 receptors in pancreatic cancer cells, respectively. (*d-f* and *j-l*) Corresponding specificity controls pre-adsorbed with the respective blocking peptide.

using an anti-FAAH antibody (Fig. 3*a*), 13 were moderately stained (Fig. 3*b*) and 13 showed a strong staining for FAAH (Fig. 3*c*). Similarly, 3 out of 31 samples were negative for MGLL (Fig. 3*g*) whereas 12 tissues were moderately positive (Fig. 3*h*) and 16 samples showed strong MGLL staining (Fig. 3*i*) in pancreatic cancer cells. Figures 3*d-3f* and 3*j-3l* show the corresponding negative controls which were incubated with the primary antibodies and the corresponding blocking peptide.

Subsequent survival analysis demonstrated that both low cancer cell FAAH and MGLL were correlated with a shorter survival of the patients (Figs. 4*a* and 4*b*; $p = 0.036$ and $p = 0.017$, respectively). The median survival of patients with low cancer cell FAAH was 10 months as compared to 19.1 months of patients with moderate-to-strong FAAH immunoreactivity. Similarly, patients with low MGLL survived in median only 8 months whereas patients with moderate-to-strong MGLL staining in pancreatic cancer cells had a median survival of 21.8 months after pancreatic cancer resection. However, these results were not confirmed in the quantitative RT-PCR showing no significant differ-

ences between FAAH/MGLL mRNA expression levels (below/above median) and survival (Figs. 4*c* and 4*d*; $p = 0.874$ and $p = 0.6$, respectively). This may be due to the low number of cancer cells in pancreatic cancer tissue specimens (which mainly consist of a fibrotic reaction—desmoplasia).

Low CB1 receptor levels in nerves are associated with strong pain symptoms

Enlarged nerves in pancreatic cancer tissues were heterogeneously stained using anti-CB1 and anti-CB2 antibodies (Figs. 5*a-5c* and 5*g-5i*) with a wide variety of unstained, moderate and strong staining patterns. Immunoreactivity of CB1 was generally more pronounced than that of CB2, while both antibodies mainly stained the perineurium of the pancreatic nerves. In some nerves with strong CB1 staining, highly positive ganglia were found (Fig. 5*c*). Consecutive sections which were incubated with the primary antibodies and the corresponding blocking peptide showed the specificity of the staining (Figs. 5*d-5f* and 5*j-5l*).




FIGURE 2 – CB1 receptor levels correlate with survival. CB receptor cancer cell protein levels as well as whole tissue mRNA expression levels were evaluated. (a) Inverse correlation ($p = 0.0011$) of high CB1 levels of pancreatic cancer cells and worse (shorter) patient survival. (b) No such correlation was observed for CB2 receptor immunostaining intensities. (c) Confirmation of the IHC quantitation results in a second, larger ($n = 53$) cohort of patients using quantitative RT-PCR: high CB1 receptor mRNA expression is related to shorter survival of the patients ($p = 0.026$). (d) CB2 receptor mRNA expression is not related to survival ($p = 0.195$).

To determine potential correlations between CB receptor levels and the extent of pain, PDAC patients were stratified according to low or high CB receptor levels in pancreatic nerves (no staining: 0; moderate staining: 1; strong staining: 2). A clinical pain score was available for 27 of 40 patients. For CB1, 18 out of 27 tissue specimens showed low receptor levels in nerves, whereas for CB2, 21 out of 27 demonstrated low receptor staining intensities. Subsequently, a correlation analysis of nonparametric data (Spearman rho) revealed an inverse correlation between CB1 and CB2 receptor levels and pain ($p = 0.041$ and $p = 0.044$, respectively). After stratification into 2 groups (low and high CB receptor levels), patients with high CB1 receptor levels had a significantly lower pain score than patients with low CB1 receptor levels ($p = 0.012$; Fig. 5m). For CB2 receptor levels, however, there was only a nonsignificant trend toward less pain in patients with high CB2 staining intensities ($p = 0.35$; Fig. 5m).

High pain scores are associated with a better outcome

Since low CB1 receptor levels were associated with increased survival but at the same time with a high pain score, we tested a potential general relationship between pain and survival in our patient collective. A log-rank test of patients with low and high pain scores revealed a statistically significant difference in survival (median survival 11 months in the group with low pain

scores versus 24.5 months in the group with high pain scores; $p = 0.0343$; Fig. 5n).

FAAH and MGLL levels on pancreatic nerves are not associated with pancreatic cancer pain

Intrapancreatic nerves in pancreatic cancer tissue samples were also heterogeneously stained when using anti-FAAH and anti-MGLL antibodies (Fig. 6): regarding FAAH, 8 samples were unstained (Fig. 6a), 14 samples were moderately (Fig. 6b) and 7 samples were strongly stained (Fig. 6c); Out of 31 samples stained with MGLL, no nerve structures were detectable in 2 samples. Out of the remaining 29 samples, 11 samples showed no immunoreactivity (Fig. 6g), 11 samples were moderately (Fig. 6h) and 7 samples were strongly stained (Fig. 6i). Figures 6d–6f and 6j–6l show the corresponding negative controls for which the primary antibodies were pre-adsorbed with the respective blocking peptide. To test whether FAAH and MGLL are associated with pain in pancreatic cancer, a quantitative evaluation of the immunoreactivities of these enzymes was carried out. This analysis revealed that there were no differences in regard to the pain score when comparing patients with no FAAH/MGLL immunoreactivity and patients with FAAH/MGLL immunopositivity on intrapancreatic nerves (Fig. 6m; $p = 0.06$ and $p = 0.582$).




FIGURE 3 – FAAH and MGLL immunostaining in pancreatic cancer. Low (*a, g*), moderate (*b, h*) and strong (*c, i*) immunostaining for FAAH or MGLL in pancreatic cancer cells, respectively. (*d–f* and *j–l*) Corresponding specificity controls pre-adsorbed with the respective blocking peptide.

Endocannabinoids AEA and AG are unchanged in PDAC

The endocannabinoids AEA and 1- and 2-arachidonoylglycerol (1 + 2-AG) were detectable in human pancreas tissues, whereas oleoylethanolamide was not found. AEA and 1 + 2-AG levels were unchanged in pancreatic cancer compared to normal human pancreas tissue samples ($p = 1.0$ and $p = 0.213$, respectively; Fig. 6*n*).

Discussion

Cannabinoids are used more frequently than earlier as supportive drugs in the palliative treatment of a variety of cancers.⁵³ In particular, their anticachectic and antiemetic properties are of clinical relevance in both chemotherapy-treated and end-stage cancer patients. The development of cannabinoid derivatives is a renaissance in medicine, since natural cannabinoids had been used by many societies until their prohibition at the beginning of the last century due to their addictive potential. However, the potential for

addiction and psychotropic side effects seems to be by far overestimated, especially for patients with malignant tumors, for whom pain control, stable weight and quality of life are the main measures of medical therapy.^{54,55}

Cannabinoids have just recently been shown to exert growth inhibitory properties in pancreatic cancer.^{16,17,23} Therefore, the most important new findings of our study are (*i*) a correlation between low cancer cell CB1 receptor or high FAAH/MGLL levels and survival, (*ii*) an inverse correlation between CB1 receptor levels on intrapancreatic nerves and the pain status of the patients and (*iii*) an inverse correlation between pain symptoms and overall survival of pancreatic cancer patients. Although there was no difference in endocannabinoid levels between healthy people and patients suffering from pancreatic cancer, cannabinoid receptors, FAAH and MGLL levels showed a significant variation. Despite heterogeneous staining intensities, levels of cannabinoid receptors were generally upregulated in pancreatic cancer suggesting an active role of these receptors in pancreatic carcinogenesis. In contrast to our findings in acute pancreatitis where there was upregu-




FIGURE 4 – FAAH and MGLL cancer cell levels correlate with survival. FAAH and MGLL cancer cell protein levels as well as whole tissue mRNA expression levels were evaluated. (a) Correlation ($p = 0.036$) of high FAAH levels of pancreatic cancer cells and longer patient survival. (b) Correlation ($p = 0.017$) of MGLL cancer cell immunostaining intensity and survival. (c, d) FAAH and MGLL mRNA whole pancreatic cancer tissue expression levels are not correlated with survival ($p = 0.874$ and $p = 0.62$, respectively).

lation of the endocannabinoid system, we did not find any changes in endocannabinoid levels in pancreatic cancer tissues. This may be due to the low number of samples tested but may also reflect the strong desmoplastic reaction in pancreatic cancer. In the vast areas of peritumoral desmoplasia, the proportion of cancer cells to stroma decreases. Therefore, although the endocannabinoid system seems not to be influenced by pancreatic cancer, one may speculate that the local concentrations of endocannabinoids in the vicinity of the tumor cells might be significantly altered. However, with the currently used techniques, such changes cannot be measured. Thus, we have to base the conclusions of our findings mainly on the status of the cannabinoid receptors and the cannabinoid metabolizing enzymes.

We found an inverse correlation between CB1 receptor levels and survival as well as a correlation between FAAH or MGLL cancer cell levels and survival. This is of particular interest since these results may partially contradict the reported antiproliferative effects of cannabinoids on pancreatic cancer growth.^{16,17} Xu *et al.* reported that overexpression of CB1 and CB2 correlated with an improved prognosis of patients with hepatocellular carcinoma.³⁶ These results point toward different roles of cannabinoid receptors in different malignancies. Furthermore, since the activity of the endocannabinoid system, its receptors and metabolizing enzyme is

a snap-shot taken during the surgical resection of the tumor, subsequent regulatory mechanisms which may occur as a consequence of a potential treatment using cannabinoids are not taken into account in our analyses. Furthermore, heterogeneous expression patterns of proteins of the cannabinoid system may preclude their potential use as clinical prediction markers. This holds particularly true since specificity and sensitivity of such a test will be strongly influenced by the respective pathologist who conducts the semi-quantitative analysis. Therefore, computerized imaging analysis systems may provide a better way to reliably quantify protein levels of these proteins for the standardized clinical application.

As can be expected based on what is known about CB receptor function in nerves,^{37,38} there was an inverse correlation between the CB1 receptor expression and pain symptoms of the patients. Similar results were obtained for CB2, although these did not reach statistical significance. Therefore, cannabinoid receptors may be involved in the development of pancreatic cancer pain. However, a lack of correlation between FAAH/MGLL levels and pain scores suggests that endocannabinoid inactivation through FAAH or MGLL seems not to be involved in pancreatic cancer pain.

Interestingly, we also observed an inverse correlation between pain symptoms and survival. This finding may be explained by the fact that pain is a warning symptom leading to earlier diagnosis of




FIGURE 5 – Cannabinoid receptor immunoreactivity of nerves in pancreatic cancer. Low (*a, g*), moderate (*b, h*) and strong (*c, i*) immunostaining for CB1 or CB2 receptors, respectively, in nerves of pancreatic cancer tissues. (*d–f* and *j–l*) Corresponding specificity controls pre-adsorbed with the respective blocking peptide. (*m*) High CB1 levels in pancreatic nerves are associated with a lower pain score ($p = 0.012$), while no such association is seen for CB2 levels ($p = 0.35$). (*n*) Kaplan–Meier survival curves of patients with low and high pain scores. Median survival was 11 months for patients with low pain scores and 24.5 months for patients with high pain scores ($p = 0.0343$).

either local recurrence of the tumor or metastasis. To analyze the value of pain as a warning symptom, we compared patients' TNM status and tumor grade with their pain scores. Our correlation analysis revealed no differences between groups with high and low levels of pain. However, both the tumor grade and the

T(N)M status are crude classifications providing little (to no) information on the microscopic tumor burden or biological behavior (other than the nodal status, which is the only prognostic factor in resectable pancreatic cancer and also in our patient collective).




FIGURE 6 – FAAH and MGLL immunoreactivity of nerves in pancreatic cancer and endocannabinoid levels. Low (*a, g*), moderate (*b, h*) and strong (*c, i*) immunostaining for FAAH or MGLL receptors, respectively, in nerves of pancreatic cancer tissues. (*d–f* and *j–l*) Corresponding negative controls. (*m*) Neither FAAH nor MGLL levels in pancreatic nerves are associated with pain ($p = 0.06$ and $p = 0.58$, respectively). (*n*) Endocannabinoid levels: 1 + 2-AG and AEA were unchanged in a comparison of normal pancreas and pancreatic cancer tissues ($p = 1.0$ and $p = 0.213$, respectively). NP: normal pancreas; PDAC: pancreatic ductal adenocarcinoma.

Taken together, our observation could—together with other reports—build the basis for a clinical testing of cannabinoids in the treatment of pancreatic cancer. Furthermore, our data strengthen the perception that cannabinoids may be useful in treating pancreatic cancer-associated pain.

Acknowledgements

The authors thank Ms. Kathrin Schneider, Ms. Monika Meinhardt and Ms. Bruni Bentzinger for excellent technical support. Authors are indebted to Ms. Judith Harvey-White and Dr. George

Kunos for the help with endocannabinoid measurements. This study was supported in part by the Intramural Research program of

National Institutes of Health/NIAAA (to P.P.) and by a postdoctoral program research grant of Heidelberg University (to C.W.M.).

References

- Kleeff J, Michalski C, Friess H, Buchler MW. Pancreatic cancer: from bench to 5-year survival. *Pancreas* 2006;33:111–8.
- Jemal A, Siegel R, Ward E, Murray T, Xu J, Smigal C, Thun MJ. Cancer statistics, 2006. *CA Cancer J Clin* 2006;56:106–30.
- Sener SF, Fremgen A, Menck HR, Winchester DP. Pancreatic cancer: a report of treatment and survival trends for 100,313 patients diagnosed from 1985–1995, using the National Cancer Database. *J Am Coll Surg* 1999;189:1–7.
- Eckel F, Schneider G, Schmid RM. Pancreatic cancer: a review of recent advances. *Expert Opin Investig Drugs* 2006;15:1395–410.
- Di Marzo V, Bifulco M, De Petrocellis L. The endocannabinoid system and its therapeutic exploitation. *Nat Rev Drug Discov* 2004;3:771–84.
- Matsuda LA, Lolait SJ, Brownstein MJ, Young AC, Bonner TI. Structure of a cannabinoid receptor and functional expression of the cloned cDNA. *Nature* 1990;346:561–4.
- Munro S, Thomas KL, Abu-Shaar M. Molecular characterization of a peripheral receptor for cannabinoids. *Nature* 1993;365:61–5.
- Giang DK, Cravatt BF. Molecular characterization of human and mouse fatty acid amide hydrolases. *Proc Natl Acad Sci USA* 1997;94:2238–42.
- Dinh TP, Carpenter D, Leslie FM, Freund TF, Katona I, Sensi SL, Kathuria S, Piomelli D. Brain monoglyceride lipase participating in endocannabinoid inactivation. *Proc Natl Acad Sci USA* 2002;99:10819–24.
- Ligresti A, Cascio MG, Di Marzo V. Endocannabinoid metabolic pathways and enzymes. *Curr Drug Targets CNS Neurol Disord* 2005;4:615–23.
- De Petrocellis L, Melck D, Palmisano A, Bisogno T, Laezza C, Bifulco M, Di Marzo V. The endogenous cannabinoid anandamide inhibits human breast cancer cell proliferation. *Proc Natl Acad Sci USA* 1998;95:8375–80.
- Guzman M. Cannabinoids: potential anticancer agents. *Nat Rev Cancer* 2003;3:745–55.
- Kogan NM. Cannabinoids and cancer. *Mini Rev Med Chem* 2005;5:941–52.
- Nithipatikom K, Endsley MP, Isbell MA, Falck JR, Iwamoto Y, Hillard CJ, Campbell WB. 2-arachidonoylglycerol: a novel inhibitor of androgen-independent prostate cancer cell invasion. *Cancer Res* 2004;64:8826–30.
- Sarfaraz S, Afaq F, Adhami VM, Mukhtar H. Cannabinoid receptor as a novel target for the treatment of prostate cancer. *Cancer Res* 2005;65:1635–41.
- Fogli S, Nieri P, Chicca A, Adinolfi B, Mariotti V, Iacopetti P, Breschi MC, Pellegrini S. Cannabinoid derivatives induce cell death in pancreatic MIA PaCa-2 cells via a receptor-independent mechanism. *FEBS Lett* 2006;580:1733–9.
- Carracedo A, Gironella M, Lorente M, Garcia S, Guzman M, Velasco G, Iovanna JL. Cannabinoids induce apoptosis of pancreatic tumor cells via endoplasmic reticulum stress-related genes. *Cancer Res* 2006;66:6748–55.
- Galve-Roperch I, Sanchez C, Cortes ML, del Pulgar TG, Izquierdo M, Guzman M. Anti-tumoral action of cannabinoids: involvement of sustained ceramide accumulation and extracellular signal-regulated kinase activation. *Nat Med* 2000;6:313–9.
- Casanova ML, Blazquez C, Martinez-Palacio J, Villanueva C, Fernandez-Acenero MJ, Huffman JW, Jorcano JL, Guzman M. Inhibition of skin tumor growth and angiogenesis in vivo by activation of cannabinoid receptors. *J Clin Invest* 2003;111:43–50.
- Grimaldi C, Pisanti S, Laezza C, Malfitano AM, Santoro A, Vitale M, Caruso MG, Notarnicola M, Iacuzzo I, Portella G, Di Marzo V, Bifulco M. Anandamide inhibits adhesion and migration of breast cancer cells. *Exp Cell Res* 2006;312:363–73.
- Vaccani A, Massi P, Colombo A, Rubino T, Parolaro D. Cannabidiol inhibits human glioma cell migration through a cannabinoid receptor-independent mechanism. *Br J Pharmacol* 2005;144:1032–6.
- Blazquez C, Casanova ML, Planas A, Del Pulgar TG, Villanueva C, Fernandez-Acenero MJ, Aragonés J, Huffman JW, Jorcano JL, Guzman M. Inhibition of tumor angiogenesis by cannabinoids. *FASEB J* 2003;17:529–31.
- Carracedo A, Lorente M, Egia A, Blazquez C, Garcia S, Giroux V, Malicet C, Villuendas R, Gironella M, Gonzalez-Feria L, Piris MA, Iovanna JL, et al. The stress-regulated protein p8 mediates cannabinoid-induced apoptosis of tumor cells. *Cancer Cell* 2006;9:301–12.
- Bifulco M, Laezza C, Valenti M, Ligresti A, Portella G, Vincenzo DIM. A new strategy to block tumor growth by inhibiting endocannabinoid inactivation. *FASEB J* 2004;18:1606–8.
- McKallip RJ, Nagarkatti M, Nagarkatti PS. Delta-9-tetrahydrocannabinol enhances breast cancer growth and metastasis by suppression of the antitumor immune response. *J Immunol* 2005;174:3281–9.
- Hart S, Fischer OM, Ullrich A. Cannabinoids induce cancer cell proliferation via tumor necrosis factor alpha-converting enzyme (TACE/ADAM17)-mediated transactivation of the epidermal growth factor receptor. *Cancer Res* 2004;64:1943–50.
- Guzman M, Duarte MJ, Blazquez C, Ravina J, Rosa MC, Galve-Roperch I, Sanchez C, Velasco G, Gonzalez-Feria L. A pilot clinical study of Delta9-tetrahydrocannabinol in patients with recurrent glioblastoma multiforme. *Br J Cancer* 2006;95:197–203.
- Wang L, Liu J, Harvey-White J, Zimmer A, Kunos G. Endocannabinoid signaling via cannabinoid receptor 1 is involved in ethanol preference and its age-dependent decline in mice. *Proc Natl Acad Sci USA* 2003;100:1393–8.
- Erkan M, Kleeff J, Esposito I, Giese T, Ketterer K, Buchler MW, Giese NA, Friess H. Loss of BNIP3 expression is a late event in pancreatic cancer contributing to chemoresistance and worsened prognosis. *Oncogene* 2005;24:4421–32.
- Michalski CW, Shi X, Reiser C, Fachinger P, Zimmermann A, Buchler MW, Di Sebastiano P, Friess H. Neurokinin-2 receptor levels correlate with intensity, frequency and duration of pain in chronic pancreatitis. *Ann Surg*, in press.
- Michalski CW, Laukert T, Sauliunaite D, Pacher P, Bergmann F, Agarwal N, Su Y, Giese T, Giese NA, Batkai S, Friess H, Kuner R. Cannabinoids ameliorate pain and reduce disease pathology in cerulein-induced acute pancreatitis. *Gastroenterology* 2007;132:1968–78.
- Michalski CW, Autschbach F, Selvaggi F, Shi X, Di Mola FF, Roggo A, Muller MW, Di Sebastiano P, Buchler MW, Giese T, Friess H. Increase in substance P precursor mRNA in noninflamed small-bowel sections in patients with Crohn's disease. *Am J Surg* 2007;193:476–81.
- Strasser F, Luftner D, Possinger K, Ernst G, Ruhstaller T, Meissner W, Ko YD, Schnelle M, Reif M, Cerny T. Comparison of orally administered cannabis extract and delta-9-tetrahydrocannabinol in treating patients with cancer-related anorexia-cachexia syndrome: a multicenter, phase III, randomized, double-blind, placebo-controlled clinical trial from the Cannabis-In-Cachexia-Study-Group. *J Clin Oncol* 2006;24:3394–400.
- Hall W, Christie M, Currow D. Cannabinoids and cancer: causation, remediation, and palliation. *Lancet Oncol* 2005;6:35–42.
- Jatoi A, Windschitl HE, Loprinzi CL, Sloan JA, Dakhil SR, Mailliard JA, Pundaleeka S, Kardinal CG, Fitch TR, Krook JE, Novotny PJ, Christensen B. Dronabinol versus megestrol acetate versus combination therapy for cancer-associated anorexia: a North Central Cancer Treatment Group study. *J Clin Oncol* 2002;20:567–73.
- Xu X, Liu Y, Huang S, Liu G, Xie C, Zhou J, Fan W, Li Q, Wang Q, Zhong D, Miao X. Overexpression of cannabinoid receptors CB1 and CB2 correlates with improved prognosis of patients with hepatocellular carcinoma. *Cancer Genet Cytogenet* 2006;171:31–8.
- Walsh D, Nelson KA, Mahmoud FA. Established and potential therapeutic applications of cannabinoids in oncology. *Support Care Cancer* 2003;11:137–43.
- Walker JM, Strangman NM, Huang SM. Cannabinoids and pain. *Pain Res Manag* 2001;6:74–9.